Pre*meno*®duo


A supportive treatment for the correction of vaginal dryness



Premeno Duo are vaginal ovules for the treatment of vaginal dryness caused by but not limited to:



- Menopause
- Stress
- Breastfeeding
- Antihormonal chemotherapy
- The use of medicines e.g. antidepressants



Premeno Duo is a non-hormonal treatment that can be used with or without local hormone therapy

- Developed in Germany **Premeno Duo** is the only dual action product on the market, healing and protecting at the same time
- Supports the healing process in damaged vaginal epithelium and reduces the risk of tears caused by mechanical effects and thus the development of bacterial infections





- Effective preventive treatment of colpitis and vaginal mycosis
- Regular use ensures adequate vaginal moistness is maintained also improving sexual sensation



Premeno Duo's active ingredients hyaluronic acid and lactic acid are an ideal, complementary combination



Hvaluronic Acid

- Promotes moisture and elasticity
- Forms a natural barrier against invading organisms
- Regulates cell development and stimulates wound healing 1,2
- Has an inhibiting effect on Candida albicans in human oral epithelial cells³



Lactic Acid / Sodium Lactate

- Restores a "healthy" pH through lactic acid substitution
- Helps prevent bacterial vaginosis^{4,5}



Premeno Duo...

- Is a handy ovule, with no requirement for gels or applicators





For every unit of Premeno Duo sold, Phoenix MedCare is donating \$1 to Sweet Louise, an organisation helping Kiwis with incurable breast cancer



[1] T.A.Dechert et al.: Hyaluronan in human acute and chronic dermal wounds, Wound Rep Reg 2006 14: 252-258

[2] J.Chen et al.: Functions of hyaluronan in wound repair: Wound Rep Reg., 1999;7:79-89

[3] A.Sakai et al.: Potential role of high molecular weight hyaluronan in the anti-Candida activity of human oral epithelial cells; Medical Mycology, 45: 73-79 [4] Mendling, Vaginose, Vaginitis, Zervizitis und Salpingitis; 2.Auflage 2006

[5] Deutsche Gesellschaft für Gynäkologie und Geburtshilfe; AWMF-Leitlinein 06/2008